

SEDIMENTATION AND DEFORMATION OF  
LOWER CRETACEOUS STRATA OF THE  
CADOMIN-LUSCAR COAL FIELD, ALBERTA

EDMONTON GEOLOGICAL SOCIETY

FALL FIELD TRIP 1988

Willem Langenberg<sup>1</sup>, Don Macdonald<sup>1</sup> and Wolfgang Kalkreuth<sup>2</sup>

<sup>1</sup> Alberta Geological Survey  
Alberta Research Council  
Edmonton, Alberta

<sup>2</sup> Institute of Sedimentary and Petroleum Geology  
Calgary, Alberta

Alberta Research Council  
Open File Report 1988-10

## CONTENTS

	Page
Preface.....	i
Acknowledgements.....	ii
Introduction.....	1
General geology.....	4
Stratigraphy and depositional environment.....	7
Structural geology.....	25
Coal quality variation.....	27
Maturation and Hydrocarbon Potential.....	34
Roadlogs and stops.....	37
References.....	61

PREFACE

The Cadomin area was the site of the first Edmonton Geological Society field trip in 1959. Subsequent trips were organized to this area by Gordon Williams in 1966 and Henry Charlesworth in 1974. After a "hiatus" of 14 years, it seems timely to go back to Cadomin. The rocks themselves have not changed, but our understanding of structural geology, sedimentology, sequence stratigraphy and ichnology has increased dramatically.

While this area has a rich variety of rock exposures ranging from Devonian to Paleocene in age, the focus of this trip is on the Lower Cretaceous part of the section. This focus arose, in part, because of the information explosion in geology over the past two decades, making it difficult to offer the older style all inclusive field trip. This field trip, therefore, reflects our sciences increased specialization.

We hope you enjoy this field trip, hang on to the guidebook and retain fond memories of your ..."fall of '88 trip to Cadomin".

## ACKNOWLEDGMENTS

We would like to acknowledge the Edmonton Geological Society for the invitation to run this trip and in particular to Mr. Donald Keith and Mr. Rudy Strobl for their assistance with logistics. The Alberta Office of Coal Research and Technology and the Alberta Research Council provided research funding for much of the basic work related to the Alberta Geological Survey's coal geology program.

Thanks are also extended to the Alberta Geological Survey for their support in logistics for this field trip. The Alberta Research Council, Graphics Department, is acknowledged for the preparation of most of the diagrams. The assistance of Ms. Shauna Treasure and Mrs. Maureen FitzGerald in helping to compile the guidebook is also very much appreciated.


## INTRODUCTION

The objective of this field trip is to examine Lower Cretaceous coal-bearing rocks of the Rocky Mountain Foothills in the Cadomin-Luscar area of central Alberta. Participants will have an opportunity to observe at first hand the lithology and complex structural relationships of the rock succession. Examples of deposition related to shallow marine, coastal plain, swamp, alluvial flood plain and alluvial channel environments will be observed and the evidence for this interpretation will be discussed. Fresh coal outcrops will be observed in the mine sites. The structure of the area will be explained with some excellent exposures and accompanying down-plunge cross sections.

The Cadomin-Luscar area is located in west-central Alberta, between latitudes 53° and 53° 8' and longitudes 117° 17' and 117° 34' (figures 1 and 2). The area forms part of the Cadomin (NTS 83F/3) and Miette (NTS 83F/4) map sheets and covers approximately 100 km<sup>2</sup>.

The existence of the thick Jewel coal seam has been known since the turn of the century. An underground mine was developed at Cadomin in 1917 and at Luscar in 1921. These mines operated until the mid 1950's. Open pit mines were opened by Cardinal River Coals Ltd. at Luscar in 1970 and Gregg River Resources Ltd. in 1983.

The Cadomin area was mapped by the Geological Survey of Canada (MacKay, 1929 and 1930). Mountjoy (1959) mapped the Miette mapsheet. The area east of Gregg River was mapped by Hill (1980). The sections along the McLeod River at Cadomin have often been used in stratigraphic studies (e.g. Mellon, 1966, 1967; McLean, 1982 and McLean and Wall, 1981). Petrographic characteristics (rank and composition) of the coals have been described by Kalkreuth (1988), Langenberg et al. (1988) and Kalkreuth and Leckie (in press).



**Figure 1.** Location of Cadomin-Luscar coalfield with pits, stratigraphic cross sections and sections.




Figure 2. Regional geology of Cadomin area, modified from MacKay (1929).

## GENERAL GEOLOGY

The Cadomin-Luscar coal field is situated in the Inner Foothills. The area is largely underlain by Lower Cretaceous rocks of the Luscar Group as defined by Langenberg and McMechan, 1985 (figures 3 and 4). The coal-bearing Luscar Group was deposited in an overall regressive sequence, during early Albian time. This sequence represents the second major, western sourced, Cretaceous clastic wedge to prograde into the Interior Cretaceous seaway. The first wedge is the Kootenay/Nikanassin succession. The Luscar Group consists of Cadomin, Gladstone, Moosebar and Gates Formations. The Cadomin Formation consists of alluvial conglomerates. The Gladstone Formation consists of alluvial sandstone, shale and minor coal and is of Aptian-Early Albian age. The Moosebar Formation contains marine shale and minor sandstone and is also of Early Albian age. The largely nonmarine Gates Formation consists of sandstones, shales and coal, and can be divided into three members, i.e., the Torrens, Grande Cache and Mountain Park Members. The age of the Gates Formation ranges from Early to Middle Albian. The Torrens Member is defined by shallow marine (shoreface) sandstones. The Grande Cache Member shows coastal- and alluvial-plain sandstones, shales and major economic coal seams. It grades into the Mountain Park Member, which consists of fluvial, fining-upward sandstones, shales, and minor coal seams. Four regional marine sedimentation cycles can be distinguished in the transition from Moosebar to Gates Formations (Macdonald et al., 1988).

Two open pit coal mines (Cardinal River Coals and Gregg River Resources) are presently producing from the Grande Cache Member. All production comes from the Jewel Seam at the base of this member and a total of 4.4 million tonnes of raw metallurgical coal was produced in 1987 (ERCB, 1987). The rank of the Jewel Seam is mainly medium volatile bituminous. The R seam (Rider Seam of Cardinal River and Ruff Seam of Gregg River) is situated about 60 m stratigraphically above the Jewel Seam. The overlying strata of the Grande Cache Member contain additional, generally thin coal seams. In addition, the Torrens, Mountain Park, Nikanassin, Cadomin and Gladstone contain thin coal seams.






Figure 3. Simplified geological map of the Cadomin-Luscar coalfield, compiled and modified from Hill (1980) and Karst (Gregg River Resources, unpublished).

		Foothills		Plains									
Age		Central Northern Alberta	Northeast British Columbia	Peace River Plains	Central Alberta	Global Cycles							
Albian	Upper	Luscarn Group Gates Torrens Mbr. Grande Cache Mbr. Mountain Park Mbr.	Moosebar Gething Bullhead Gp.	Spirit River Wilrich Mbr. Falher Member Notikewin Member	Paddy Mbr.	Viking	UZA-2						
					Joli Fou	UZA-1 1.5 ? 1.4 ? 1.3 ? 1.2 ? 1.1							
	Cadotte Mbr.				Boulder Ck.		Mannville Group Glauc. Mbr.						
	Harmon Mbr.				Hulcross			Ostracode Mbr.					
	Lower				Luscarn Group		Gates	Moosebar	Spirit River	Mannville Group	Glauc. Mbr.	LZB-4	
													Grand Rapids
													Clearwater
													Deville
	Aptian				Luscarn Group		Gladstone	Gething	Gething	Bluesky Fm.	Ostracode Mbr.	LZB-4	
						Elleslie or Basal Quartz							
?	Luscarn Group	Cadomin	Cadomin	Cad.	Jurassic - Devonian	Miss - Devonian							
								Nikanassin	Minnes Gp.				

Figure 4. Lower Cretaceous stratigraphic nomenclature of northcentral Alberta and northeastern British Columbia.

The rocks of the area are highly deformed by folding and faulting, with major structures such as the Cadomin Syncline, the Luscar Anticline and the Drinnan Thrust (figures 5 and 6). In the south, the area is bounded by the Nikanassin Thrust, which forms the boundary between Foot-hills and Mountains in the region.

## STRATIGRAPHY AND DEPOSITIONAL ENVIRONMENTS

### NIKANASSIN FORMATION

The Nikanassin Formation straddles the Jurassic/Cretaceous boundary and is thought to represent the first western sourced clastics into the West Alberta foreland basin (Stott, 1984). The formation is generally divisible into a lower marine and an upper coastal and alluvial plain portion. The correlative Kootenay Formation, in central and southern Alberta, is coal-bearing and represents a more complete coastal plain to alluvial plain environment. At Cadomin, the upper Nikanassin is exposed immediately below the Cadomin Formation, along the McLeod River railroad section. Some gas is found in the Nikanassin in the Elsworth Deep Basin.

### LUSCAR GROUP

The Luscar Group consists of the Cadomin, Gladstone, Moosebar and Gates Formations. This group, which is equivalent to the Mannville Group of central Alberta (figure 4), shows both marine and non-marine sedimentary environments. In the Deep Basin (north of the Cadomin area), as many as eight marine cycles of marine sedimentation have been recognized in correlative strata of the Spirit River Formation (Cant, 1983; Smith et al; 1984). These include the Wilrich A and B cycles, the Falher A through G cycles and the Notikewin (figure 7). In the western deformed part of the Deep Basin (Cadomin area), the Gates Formation is mostly nonmarine and it is not possible to recognize all of the Falher Member cycles (figure 7).







Figure 6. Cross sections XX' and YY' through the Cadomin-Luscar coalfield. Section XX' is on the Gregg River property (after Karst, Gregg River Resources Ltd., unpublished). Section YY' is on the Cardinal River property (modified from Hill, 1980).









- |   |   |   |  |
|---|---|---|--|
|  | Continental fluvial                         |  | Coal seam, carbonaceous shale<br>(thickness exaggerated) |
|  | Continental alluvial plain                  | J   | Jewel seam   |
|  | Brackish                                    | R   | Rider seam   |
|  | Marine, nearshore                           | Not.  | Notikewin  |
|  | Marine, offshore, transgressive<br>deposits | 3, 4, etc.  | Grande Cache coal seam nomenclature                      |
| .....   | Sequence boundary (approximate)             | A, B, etc.  | Father cycles  |

Figure 7. Schematic cross section of the Gates Formation clastic wedge, Cadomin to Elmworth area.

## CADOMIN FORMATION

The Cadomin Formation was deposited in alluvial fans and on braided river pediment plains with chert pebble conglomerates, and is unconformably overlying the Nikanassin (McLean 1977, Gies, 1984). Much of the Cadomin that developed in the west as alluvial fans was transported and reworked into the northwest trending Spirit River channel that developed parallel to the foreland basin during the early Aptian. The Cadomin likely represents a major sea level lowering event and possibly a major sequence boundary.

Large reserves of gas are present in the Cadomin Formation in the northern Deep Basin region. At least one gas well, 10 km east of Cadomin, encountered gas in the Cadomin Formation. The Cadomin Formation is an excellent marker horizon, both at the surface and in the subsurface.

## GLADSTONE FORMATION

The Lower Cretaceous (Aptian-Albian) Gladstone Formation is equivalent to the coal-bearing Gething Formation in northeastern British Columbia and the oil sands-bearing McMurray Formation in northeastern Alberta. The formation lies conformably on the Cadomin with no apparent major stratigraphic break. The section at Cadomin shows a transition from well-drained alluvial plain deposits with thin coals near the base, to coastal plain deposits with thicker coals near the top.


The Bluesky Member occurs at the top of the Gladstone as a near-shore marine transgressive, upward-coarsening sequence. Gas is produced from the northeast end of a barrier island or bar sand in the Bluesky in the Edson gas field. This bar sand extends to within 20 km of the Cadomin area (Jackson, 1984), although it is not gas-bearing in this region.

## MOOSEBAR FORMATION

Macdonald, et al. (1988) suggests that the marine strata of the Moosebar/Gates transition at Cadomin are likely correlative with the subsurface Wilrich and lower Falher cycles (figure 7). At Cadomin, these strata are divided into the 1st, 2nd and 3rd regional cycles (figure 8), forming a series of prograding shorelines and coastal plain deposits. A 4th cycle is present above the Jewel coal seam and is thought to represent a brackish water facies. More local scale cycles are also present in the outcrop area, probably representing local facies changes or subsidence events. The fully marine succession includes offshore to lower shoreface (with storm deposits) and shoreface to foreshore (and possibly beach, figure 8). It is uncertain, at this time, what type of prograding shorelines are represented in the Cadomin area. Very little evidence is present to support a tide or river dominated delta setting from sedimentological observations in the sections studied. McLean and Wall (1981) suggest a wave influenced strandline environment for the Cadomin area. In the subsurface to the east of Cadomin, Jackson (1984) suggests that southwest-northeast trending wave dominated shorelines were predominant during this time. There is general agreement amongst all workers that paleoflow was towards the north and Taylor and Walker (1981) argue specifically for a northwest paleoflow, parallel to the Cordillera, in the Nordegg area.

The lower Moosebar Formation (lower Wilrich or 1st cycle) in this area consists of a series of fine-grained mudstones interbedded with sharp based siltstones and thin sandstones (figure 8). A thin argillaceous coal is present, as well as *Planolites* and *Skolithos* burrows. The cycle terminates in a 3 m coarsening-up sequence that is capped by a thin rooted coal. Well preserved conifer cones and leaf impressions are present in a thin shale bed near the coal. This cycle is best exposed on the Cardinal River Coal property (50A3 pit), near the southwestern boundary of the Cadomin-Luscar coalfield (figure 1).





Legend:

- |   |   |                           |                         |
|---|---|---------------------------|-------------------------|
| Low-angle x-strata                      | Carbonaceous matter in thin laminae                     | Ironstone band            | Basal lag               |
| Hummocky cross stratification           | Tectonically thinned coal                               | Erosional contact         | Parallel stratification |
| Very fn interlaminated                  | Fault   | Brackish                  | Intraclasts             |
| Carbonaceous matter-particles           | Glaucinitic   | Trough x-strata           | Graded bedding          |
| Shells (marine)                         | Bioturbated   | Ripples                   | Slickensides            |
| Fining-upward cycle                     | Forams  | Soft sediment deformation |                         |
| Coarsening-upward cycle                 | Gradational contact                                     | Roots                     |                         |
| Carbonaceous matter-finely disseminated | Sharp contact   | Leaf imprints             |                         |
|   | Burrows (vertical, horizontal, with spriten, branching) | Log or stems              |                         |

Figure 8. Cross section showing the marine to nonmarine transition, Cadomin-Luscar area.

This first cycle in the Moosebar Formation is interpreted to have formed in offshore to shoreface environments in which storm events periodically deposited thin sand units. In this area, the cycle terminated in an emergent bar. A fairly major relative lowering of sea level, and consequent exposing of the shelf, is indicated by the preservation of well preserved plant fossils and the development of thin rooted coals.

The lower part of the 2nd marine cycle corresponds to the upper part of the Moosebar Formation. A glauconitic, sharp based pebble conglomerate bed, commonly graded, is found near the base of the cycle in Section 6 and 9 (figure 8) and is interpreted as an offshore transgressive deposit. It may, in part, represent very slow deposition of a condensed section as described by Haq et al. (1987). The presence of glauconite in the pebble beds supports this idea (Weimer, 1983). The boundary between Moosebar and Torrens is gradational and is somewhere in the 2nd cycle. Alternatively, it could be placed either at the top or the base of the 2nd cycle.

#### GATES FORMATION

The Gates Formation consists of the Torrens, Grande Cache and Mountain Park Members. Both shallow marine and non-marine sedimentary environments are deduced for this formation.

#### TORRENS MEMBER

The base of the Torrens Member is not well defined, but is somewhere in the 2nd marine cycle. The 2nd cycle is believed to correspond to one of the cycles in the Wilrich Member (figure 8). This cycle, at Cadomin, consists of two or more coarsening-up sequences stacked on top of each other. The coarsening-up cycles are commonly heterolithic, with fine sand dominating. Hummocky cross stratification, soft sediment deformation, wave ripples, and parallel laminations are all found in this cycle. Trace fossils include *Planolites*, *Diplocraterion* and *Skolithos*. The second cycle is capped by a rooted

coal or carbonaceous mudstone, herein called the "Torrens A-coal". This coal can be traced in the subsurface as far north as township 57 (south of Grande Cache), where it disappears in another coarsening-up sequence (Macdonald et al., 1988). The Torrens A-coal is seen to vary from a 90 cm coal in the south (Section 9) to a carbonaceous shale at Section 6, and is locally eroded at Section 5 (figure 8).

The second cycle is interpreted to have formed primarily in the offshore to shoreface environments. The offshore/transgressive deposits of the Moosebar give way upsection to lower shoreface and finally foreshore sediments of the Torrens Member. The trace fossil assemblage is consistent with this interpretation. The hummocky cross stratification supports the storm deposited origin and has been found in other locations of the Moosebar to Gates transition (Leckie and Walker, 1982). The laterally extensive Torrens A-coal in the subsurface may suggest sea level lowering at this time (Macdonald et al., 1988).

The 3rd marine cycle consists of massive (though occasionally thinly bedded), fine to very fine-grained sandstone, which has been traditionally been called the Torrens Member. Faint parallel laminations are the predominant structures, with some trough cross bedding and hummocky cross stratification also present (figure 6). Scour surfaces, with pebble lag deposits, are also common in the cycle. Mudstone is a very minor lithology in this cycle and is usually associated with the hummocky cross stratification. Trace fossils are particularly abundant near the top of the 3rd cycle and include *Skolithos*, *Ophiomorpha*, *Planolites*, and *Diplocraterion*. The sequence is capped by the thick Jewel Seam. The sandstones below the coal seam are commonly well rooted and show a discoloration in the top 50 cm. Tree trunk impressions and root casts can be found on the uppermost bedding plane surface of these sandstones. A more medium-grained, intraclast rich, trough cross stratified facies occurs at one locality within this area.

The 3rd cycle is probably correlative with the Falher G or Wilrich A cycles in the Deep Basin (Macdonald et al., 1988). This cycle is

interpreted to be a succession of two prograding shorelines sequences, with upper shoreface to foreshore environment transitions being present. The thin Torrens "B-coal" that is present at Section 9, very likely represents a shifting away from this area of the prevailing sediment source and the development of a small lagoonal environment. This coal is not traceable throughout the Cadomin area (figure 8). The trace fossil assemblage is typical of the *Skolithos* ichnofacies, which is most commonly found in high energy beach or beach-like facies (Ekdale et al., 1984). The presence of hummocky cross stratification high up in the sequence precludes a true beach environment for parts of the Torrens, because these structures generally form below fair wave base (Walker, 1984). The more medium-grained, trough cross stratified facies is interpreted to be a later tidal channel deposit.

#### LOWER GRANDE CACHE MEMBER

The interval between the Jewel and R seams can be divided into five lithofacies units (figure 9). Lithofacies GC1 is the 10 m thick Jewel coal seam which overlies the Torrens Member, has a well rooted seatearth, which often contains log and root impressions (figure 8). The Jewel Seam forms the base of the Grande Cache Member. Macdonald et al. (1988) suggests that part of the Jewel Seam is stratigraphically equivalent, though not necessarily time equivalent, to the No. 3 seam at Grande Cache. To the east, in the subsurface, the Jewel Seam appears to be correlative with the Mannville "Medicine River coal marker" described by Strobl (1988). Farther towards the north, into the Deep Basin, the Jewel equivalent is seen to disappear into marine strata (figure 7). This evidence suggests a more landward position for the peat swamp development, perhaps up to 100 km from the coastline on a coastal plain.

Englund et al. (1986) describe low ash, low sulfur, Pennsylvanian coastal coals, having no overlying marine strata, that are believed to be directly related to the development of peat swamps on top of abandoned delta lobes in an overall regressive sequence. McCabe (1987) describes coals from the Upper Cretaceous Horseshoe Canyon Formation and Judith River Group, forming in north-south belts parallel to the




Figure 9. Stratigraphic cross section showing facies interpretation for the Grande Cache Member between the Jewel and R seams.

paleocoastlines and having the thickest accumulation of peats (coals) some 40-80 km landward of the shoreline. The Okefenokee Swamp is cited as a modern day analog for the Upper Cretaceous coals by McCabe (1987), and this model may equally well apply to the Lower Cretaceous Jewel Seam. McCabe (1984) has suggested that in order for thick, low ash coals to form in a low lying coastal plain swamp, they must be some distance landward from active shoreline processes, yet still maintain high water tables. This would seem to be the case for the Jewel Seam at Cadomin. Because of the close connection with regionally extensive foreshore deposits below the coal, Kalkreuth and Leckie (in press, their figures 1 and 19) introduced the term strandplain coals for these coastal plain coals.

A 4th marine cycle (directly above the Jewel Seam) is designated as lithofacies unit GC2 and extends upsection until the first major fluvial sandstones are encountered (figure 9). A brackish water interpretation is based on the presence of trace fossils (*Diplocraterion* and *Planolites*) at Section 6, and the presence of lenticular bedding throughout this interval. Specimens of the siliceous forams *Hippocrepina* (?) sp., *Miliammina* (?) sp. and *Saccamina* sp. have been recovered from three locations above the Jewel Seam in this area and are indicative of ... "a shallow brackish (not normal marine) marine environment" (Wall, pers. comm.). The fourth cycle is thought to be related to the transgressive portion of the Falher F cycle in the Deep Basin further north.

The nonmarine sequence above the tidal flat facies GC2 (figure 9) is non coal-bearing or contains only very thin seams. No trace or hard-bodied marine fossils have been observed above the tidal flat facies, at least up to the R coal seam.

The GC3 lithofacies unit overlies the tidal flat unit and consists of thick, amalgamated, trough-cross stratified, log-bearing, fine-grained sandstones (figure 9). Some large lateral accretion bedding can be seen in the 51B2 pit at the Cardinal River minesite. The unit is interpreted to be a series of amalgamated fluvial channel deposits. A

meandering system is suggested by the lateral accretion bedding.

The GC4 lithofacies consists of a series of thin, fining-upward sandstones and mudstones and some coarsening-upward sequences. A very thin (10 - 20 cm) rooted coal caps this unit over most areas, except where channeling has removed the seam (figure 7). The GC4 lithofacies is interpreted to be a series of stacked, small fluvial channels, with associated crevasse splay deposits.

Over most of the area the Unit GC5 consists of fine-grained mudstones, thin discontinuous sandstones, the R seam capping the unit and minor coarsening-up sequences. Section 4 (figure 9), in contrast, shows a series of fining up cycles, each about 2 - 6 m thick. This unit is interpreted to be primarily a floodplain deposit, with one area of stacked, small fluvial channel deposits. Paleocurrent measurements are not available, but the cross-section F-F' orientation (figure 9), suggests either an east-west or northwest-southeast orientation for this channel system. The R seam is considered to be a typical alluvial plain coal, which was probably deposited contemporaneously with very small fluvial channels. It is possible, however, that the R seam is related to one of the Falher cycles (B or C?) during the post regression, coal forming period. The R seam does lie at about the same stratigraphic position as the #10 or 11 seam at Grande Cache (figure 6), both of which are thought to be associated with the Falher cycles in the Elsworth area (Macdonald et al., 1988 ).

#### UPPER GRANDE CACHE MEMBER

The upper Grande Cache Member sequence above the R seam is much the same as seen in the lower portion. No significant change occurs in the interpreted depositional environment until the Mountain Park Member is reached.

#### MOUNTAIN PARK MEMBER

The Mountain Park is generally defined as .."the first major

greenish colored sandstone encountered going upsection" (McLean, 1977) for the purposes of field mapping in this area. These sandstones are interpreted to be large scale fluvial deposits and may occur at various stratigraphic levels, making the field mapping criteria somewhat arbitrary. The uppermost coal seam (>0.6 m thick) in the Grande Cache Member is alternately used, when no major greenish sandstone is present.

The contact between the Mountain Park Member and the overlying dark grey mudstones of the Blackstone Formation (Shaftesbury) is generally sharp and usually contains a thin transgressive pebble lag deposit. Thin upward-coarsening cycles are occasionally seen in the lower part of the Blackstone Formation. The Blackstone transgression may be related to the accretion of terrains during the Columbian Orogeny (Stott, 1984 and Monger, 1984). The Blackstone Formation may also be an important source rock for gas generation in this area.

#### COAL DEPOSITIONAL ENVIRONMENTS

Facies-critical macerals and a number of petrographic indices can be used to define coal facies and environments of coal deposition (Diessel, 1986). Macerals are defined as the microscopically recognizable constituents in coals. The term maceral was introduced as an analogy to the term mineral of inorganic rocks. The macerals are commonly classified in three groups: vitrinite, inertinite and liptinite. The relationships of coal macerals and depositional settings of the Lower Cretaceous Gates coals have been examined recently (Kalkreuth and Leckie, in press).

The proportions of the macerals in coals reflect the various organic source materials contributing to the accumulation of the ancient peat and the conditions prevailing during accumulation at or near surface; i.e. height of water table, pH, decay by aerobic and anaerobic bacteria, etc. Facies-critical macerals are components such as vitrinite A, which indicates on origin from wood-producing plants and liptinite macerals such as sporinite and cutinite, which refer to specific precursor materials. Others like alginite, semi-fusinite and



fusinite are indicators of the relative position of the water table during peat accumulation. Petrographic indices such as the Tissue Preservation Index and the Gelification Index (Diessel, 1986) are used to assess the depositional environments, i.e. moor types, height of water tables, preservation of cell-tissues, degree of oxidation and degradation due to transport mechanism (the Tissue Preservation Index, TPI, and Gelification Index, GI, are explained in figure 10).

Petrographic characteristics and coal depositional environments will be discussed in detail for a section from Gregg River's PQ and LM pits (see also Kalkreuth and Leckie, in press). The section sampled is shown in figure 10 and represents a composite of two sections separated by 650 m. Samples were collected from the Torrens A coal seam in between shoreface sandstones, the Jewel Seam directly on top of the shoreface sandstones and the R Seam (Ruff) from the non-marine environments (figure 9). Within this sequence, systematic trends in maceral contents, mineral matter content and in the petrographic indices are apparent (figure 10). The Torrens A Seam has a very high vitrinite content (93 vol. %) while inertinite content is low (7 vol. %). Liptinite macerals were found only in traces. The mineral matter content is very high (47 vol. %). The samples collected from the Jewel Seam show in contrast a drastic decrease in vitrinite contents (46 - 58 vol. %) while inertinite macerals account for 40 - 52 vol. %. Macerals of the liptinite group (sporinite) are still rare with up to 2 vol. % only. Mineral matter contents in the Jewel Seam range from 2 - 23 vol. %. A return to a higher vitrinite content is indicated for the Ruff Seam (96 vol. %) with only minor contributions of inertinite (3 vol. %) and liptinite (1 vol. %). The Ruff Seam has at this location a mineral matter content of 21 vol. %. Within the group of petrographic indices many show distinct changes in respect of the stratigraphic position of the coal seams. Related to the overall contents of vitrinite and inertinite macerals, the Jewel Seam has a very low V/I (vitrinite/inertinite) ratio while the Torrens A and the Ruff Seams have high ratios. Preservation of the organic materials appears to be best in the Ruff Seam which is characterized by a high vitrinite A/vitrinite B ratio, and a very high Tissue Preservation Index. In contrast, the




Figure 10. Maceral data, mineral matter contents and petrographic indices for Lower Cretaceous Gates coals, Gregg River Mine, Luscar (from Kalkreuth and Leckie, in press). Petrographic indices are: VA/VB ratio = Vitrinite A / Vitrinite B; SF/F ratio = Semifusinite / Fusinite; V/I ratio = Vitrinite / Inertinite; T/F ratio = Vitrinite / (Fusinite + Semifusinite); IR ratio = (Semifusinite + Fusinite) / (Inertodetrinite + Macrinite + Micrinite); W/D ratio = (Vitrinite A + Fusinite + Semifusinite) / (Alginite + Sporinite + Inertodetrinite); TPI ratio = (Vitrinite A + Fusinite + Semifusinite) / (Vitrinite B + Macrinite + Inertodetrinite); GI ratio = (Vitrinite + Macrinite) / (Semifusinite + Fusinite + Inertodetrinite); S/D ratio = (Vitrinite A + Fusinite + Semifusinite) / (Alginite + Sporinite + Inertodetrinite + Vitrinite B + Vitrodetrinite); Rmax% = Mean maximum vitrinite reflectance.

Jewel Seam appears to have a lesser degree of plant preservation, mainly because of substantial amounts of inertodetrinite and vitrinite B (low VA/VB ratio, figure 10). Within the inertinite group, semifusinite is the predominant maceral (22 - 24 vol. %, high SF/F ratios, figure 10) while the total amount of structured inertinite (fusinite and semifusinite) is always greater than inertodetrinite (IR ratios = 1.35 - 2.02, figure 10).

To assess the type of prevailing mire types during accumulation of the organic matter, the Gelification and Tissue Preservation Indices for the five coals have been plotted in a facies diagram (figure 11), as proposed by Diessel (1986). The three samples of the Jewel Seam plot very close to each other, indicating a similar type of paleo-environment of deposition for these coastal plain coals (strandplain coals of Kalkreuth and Leckie, in press). They must have been formed under relatively dry conditions as indicated by the high amounts of inertinite macerals (semifusinite, inertodetrinite, to a lesser extent fusinite). The major split of the seam, as illustrated in figure 10, might be a result of local flooding adjacent to fluvial/tidal channels or a result of lacustrine conditions caused by temporarily and locally increased subsidence rates. The Torrens A Seam has a much higher Gelification Index (figure 10) due to a very high vitrinite content and low amounts




Figure 11. Facies diagram for Lower Cretaceous coals at Gregg River Mine, Luscar (from Kalkreuth and Leckie, in press).

of inertinite. The formation of the seam most likely reflects slight sea level fluctuations during deposition of the Torrens Member in which peat accumulated while a relatively high water table level was maintained as indicated by the large amounts of gelified plant remains. The Ruff Seam in the Gregg River Mine area is highly variable in thickness, ash content and lateral continuity. The position in figure 11 shows very high Gelification and Tissue Preservation Indices due to the fact that the coal is made up almost entirely of vitrinite. As such, the Ruff Seam might represent a locally flooded mire of the upper delta plain or alluvial plain with a substantial input of mineral matter associated with a relatively high water table.

Petrographic indices of 16 Jewel Seam samples from the Cadomin-Luscar area (including the seams discussed previously) show that these coals form distinct groups, although considerable petrographic variation exists. In addition, some overlap of petrographic characteristics occur with the overlying coals from the non-marine environments. The V/I ratios range from 0.90 - 2.10 (mean = 1.39) and indicate, for most of the coals, the predominance of gelified components (vitrinite) over non-gelified (inerts). Within the vitrinite group the VA/VB ratio indicates a dominance of vitrinite B over vitrinite A (mean = 0.87) suggesting that substantial amounts of woody materials were decomposed prior to final deposition. The same holds true for the relatively high proportion of inertodetrinite derived from mechanical breakdown of fusinite and semifusinite precursors. The IR ratios (semifusinite + fusinite/inertodetrinite) of the Jewel Seam range from 1.20 - 2.30 with a mean of 1.58. That means that in all of these coals the sum of semifusinite + fusinite macerals is greater than inertodetrinite. IR ratios <2 were considered by Diessel (1982) to reflect hypautochthonous and allochthonous conditions in the ancient peat swamps where redeposited semifusinite, fusinite and inertodetrinite would make up the bulk of the inerts. In other words, some transportation of the peat has taken place in the course of the accumulation of the Lower Cretaceous coastal plain coals.

Tissue Preservation Indices, such as T/F ratio and TPI ratio, are

low to intermediate with values ranging from 0.50 - 1.60 for the T/F ratio (mean = 1.02) and from 0.80 - 1.54 for the TPI (mean = 1.12). These values indicate for the Jewel Seam a forest type swamp depositional environment with periods of low water tables during which substantial amounts of oxidized and partly oxidized materials were formed. The relatively high contents in the non-gelified macerals semifusinite, fusinite and inertodetrinite in respect to the gelified components vitrinite and macrinite leads to relatively low gelification indices ranging from 1.00 - 2.30 (mean = 1.55).

From the coal facies diagram (Diessel, 1986), a forest swamp environment of deposition is deduced for many of the coastal plain coals of the Lower Cretaceous (Kalkreuth and Leckie, in press). For some coals, characterized by somewhat higher gelification indices (GI) while retaining similar tissue preservation indices (TPI), a shift to less forested fen-like depositional environments is indicated with a larger input of aquatic plants such as reeds and sedges.

#### STRUCTURAL GEOLOGY

The structure of the area is shown on the accompanying geological maps (figures 2 and 3) and cross sections (figures 5 and 6). The main structures in the Cadomin area are the Cadomin Syncline, Luscar Anticline, Drinnan Thrust and Nikanassin Thrust.

#### FOLDING

Many of the macroscopic folds have relatively straight limbs and short hinge areas, and could be classified as chevron folds. Folding is cylindrical on a local scale. This means that the folds can be described by the movement of a line (which is the fold axis) parallel to itself. This implies that a cylindrical fold does not change geometry in the direction of the fold axis. However, on a regional scale folds change geometry in the direction of the fold axis and consequently they have to be considered non-cylindrical. This non-cylindricity resulted in the necessity to divide the Cardinal River property into 45 statist-

ically cylindrical domains (Hill, 1980). These cylindrical domains enabled downplunge cross sections to be constructed. Cross section YY' (figure 6) is based on a composite downplunge projection of many cylindrical domains, similar to the cross sections constructed by Langenberg et al. (1987) in the Grande Cache area. This cross section shows overturned folds, which are generally open with interlimb angles of 120 to 40°. However, several of the folds are tight with interlimb angles less than 30°. The shortening can be measured to be around 50%, using the Jewel Seam as reference horizon.

The overturned southwest limb of the Cadomin Syncline is well exposed in Gregg River's PQ Pit, in Cardinal River's 50A3 and 50A5 Pits and along the McLeod River. The plunge of this fold is generally to the southeast, but several subsidiary folds plunging to the northwest indicate the non-cylindrical nature of the folding on a regional scale. Southeasterly plunging subsidiary folds are present in Gregg River's PQ and LM pits and Cardinal River's 51A3 Pit, while northwesterly plunging folds are found in Gregg River's LM pits and Cardinal River's 50A5 Pit.

A good cross section of the Luscar Anticline is provided by the 51B3 Pit. This exposure shows that folding has had a profound effect on coal quality by the resulting thickening in fold hinges. This process was described for the Grande Cache area by Langenberg et al. (1987). Most open pits are along the hinges of folds, indicating the economic significance of the thickening. Coal has flowed into these hinges from the nearby limbs. As a result, the folding of the Torrens sandstone is more open than the strata overlying the Jewel Seam (disharmonious folding). The plunge of the Luscar Anticline is generally to the southeast.

#### FAULTING

The Nikanassin Thrust is a major thrust fault, which in places thrusts Devonian strata (Palliser Formation) over Jurassic/Cretaceous strata (Nikanassin Formation). It forms the boundary between Foothills and Mountains in the region. The Drinnan Thrust, which is well exposed

in Gregg River's PQ Pit, thrusts Nikanassin Formation over the Grande Cache Member with about 500 m of displacement. The Mystery Lake Thrust can be followed from Mountjoy's (1959) mapping in the Miette mapsheet to just west of a prominent syncline (figure 3). Consequently the fault has terminated and the shortening is transferred to the syncline. Coal in this syncline has been mined in Gregg River's AB Pit. Alternatively the fault in the southwesterly limb of the Luscar Anticline (named Luscar Thrust by Hill, 1980) could be an extension of the Mystery Lake Thrust. An unnamed fault places Nikanassin Formation over the Gladstone Formation along the Gregg River, north of the Mystery Lake Thrust. Evidence for the presence of the Folding Mountain Thrust, mapped by Mountjoy (1959) in the northern part of the study area, is lacking at the surface.

Most of the faults are southwest dipping thrust faults, however in several of the pits, northeast dipping faults with hanging-wall moved down can be observed. Hill (1980) suggested that these faults were pre-folding thrust faults. However, because these faults are not folded, they probably formed in a late stage of the folding process or post-folding. They might be a type of normal faults.

Another important structure is the Blackstone zone of detachment (figure 5), which is situated in the Blackstone Formation in between the disharmonious structures of the Cardium and Gates Formations (Hill, 1980).

#### COAL QUALITY VARIATION

Coal quality variations have been discussed in detail by Langenberg et al. (1988). Coal quality was mainly determined by proximate and ultimate analyses and petrographic analyses using incident light microscopy.

In the context of this guidebook, coal quality variations will be discussed briefly in terms of ash and volatile matter contents obtained from proximate analyses, sulfur contents based on ultimate analyses, and

vitroinite reflectance and maceral content based on petrographic analyses. Ash and sulfur contents and the variation in organic constituents (macerals) are largely determined by the original sedimentary environment, while volatile matter content and vitroinite reflectances are largely related to subsequent burial of the organic matter and to some extent to deformation.

#### ASH

Ash in the Jewel Seam comes either from visible rock partings or from finely disseminated mineral matter within the coal. In places the disseminated ash content may have increased by tectonic shearing. Qualitative field observations suggest that the more highly sheared a coal is, the more likely it is to have elevated ash values. These factors make ash percentage prediction difficult. Average disseminated ash contents of the Jewel Seam is about 14%, although in-seam ash profiles show large variations from this average (figure 12). Vertical successions with upward increase in ash and low ash zones through the center of the seam can be explained by the original acidity of the swamp (Langenberg et al., 1988).

The overall depositional setting of the Jewel Seam is believed to have been on a coastal plain, perhaps up to 100 km inland from the shorelines. Thus the Jewel Seam was allowed to attain its large thickness well removed from marine clastic influences. Smaller fluvial or tidal channels may have dissected or been adjacent to the Jewel swamp providing the visible partings. For mining purposes, a 2 m sampling interval of the Jewel Seam results in a good characterization of ash variation.

Ash values in R, Torrens and Moosebar coals all tend to be high, and are more likely related to nearby clastic environments than to the original acidity of the swamp.






Figure 12. Stratigraphic cross section E-E' showing pit-scale in-seam ash variations in the Jewel seam, throughout the Cardinal River Coals Ltd. 51B3 pit.

## VOLATILE MATTER

Volatile matter (dry and ash free) of the Jewel Seam ranges from 19.3 to 39.8%, where the higher percentages represent oxidized coal. This range indicates largely medium volatile bituminous rank. There is a systematic variation of volatile matter in the study area, where the central part has the lowest percentages (see section on vitrinite reflectance).

## SULFUR

Sulfur contents of the Jewel Seam are low compared to many other coal deposits and average 0.3%. Low sulfur combined with moderate high ash and high inertinite contents point to acidic swamps, with intermittent aerobic conditions (Cecil et al., 1980). Sulfur, which is mostly organic in the Jewel Seam, often shows elevated values at the base, and to some extent at the top of the seam. Basal elevated sulfur values can drop back to average values over short distances. Elevated sulfur values at the top of the seam may be related to overlying thick channel or brackish water deposits. A very low sulfur zone (i.e. <0.2% dry) is often found in the middle of the Jewel Seam, particularly when there are few partings.

Sulfur values in the R Seam (alluvial plain setting) generally lie around 0.4%. Sulfur values in coals found in the Torrens Member and Moosebar Formation (marine to transitional environment) tend to be higher than the Jewel Seam (i.e. >0.7% dry) and support the general model of high sulfur coals associated with overlying marine strata.

## MACERALS

Macerals (a term which was introduced as an analogy to minerals of inorganic rocks) are defined as the microscopically recognizable constituents in coals. The macerals are commonly classified in three groups: vitrinite, inertinite and liptinite. The proportions of the macerals in coals reflect the various organic source materials

contributing to the accumulation of the ancient peat and the conditions prevailing during accumulation at or near surface; i.e. height of water table, pH, decay by aerobic and anaerobic bacteria, etc. The determination of macerals in coals is complicated by the fact that many undergo severe alterations during the course of coalification. For details the reader is referred to Stach (1982) and Bustin et al. (1983).

The Jewel Seam of the Cadomin-Luscar area is characterized by relatively low vitrinite contents ranging from 46 - 67 vol % with a mean of 56 vol % (based on 16 composite samples). In contrast inertinite contents are relatively high ranging from 32 - 52 vol % with a mean of 43 vol %. Within the inertinite group major components are semifusinite (range = 17 - 24 vol %, mean = 21 vol %), inertodetrinite (range = 8 - 19 vol %, mean = 14 vol %) and fusinite (range = 3 - 7 vol %, mean = 5 vol %). Liptinite content is in general low, ranging from nil to 2 vol % with an average of 1 vol %. The relatively low vitrinite contents and high amounts of inertinite macerals, in particular semifusinite and fusinite indicate rather low water tables during peat accumulation in which a substantial part of the organic matter was oxidized prior to final sedimentation. The optically determined mineral matter is in most coals low to moderate, ranging from 2 - 23 vol % with a mean of 9 vol %. These maceral contents are very similar to those of coal seams believed to have accumulated under similar environments to the north in the Grande Cache area, Alberta and Tumbler Ridge area, B.C. (see Kalkreuth and Leckie, in press). Petrographic indices, based on these maceral contents, were presented in the section on "coal depositional environments".

#### VITRINITE REFLECTANCE

Maximum vitrinite reflectance for the Jewel Seam ranges from 0.97 to 1.43% (figure 13). The highest rank is found in the central part of the study area, along the southwest limb of the Luscar Anticline. High volatile bituminous coals are present in the southeast. The rank variation indicates syndeformational coalification, although it cannot be excluded that the central part of the coal field was buried somewhat




Figure 13. Regional vitrinite reflectance variations of the Jewel Seam.

deeper, resulting in the higher rank. However, the burial during sedimentation was probably also a result of deformation, which was taking place further west during the existence of the foreland basin. It should also be noticed that the highest ranks are not along the axis of the Cadomin Syncline as would be expected in the syndeformational coalification model, but along the southwestern flanks of the Luscar Anticline (figure 3). This might be explained by a late stage adjustment of the structure, where parts of the Cadomin Syncline have moved upwards by secondary folding and thrusting (Mystery Lake Fault and fault near Luscar).

A good linear correlation between maximum vitrinite reflectance and volatile matter (dry and ash free) is observed, enabling volatile matter to be determined from vitrinite reflectance. The map of maximum vitrinite reflectance variation (figure 13) can be used to predict rank and volatile matter of the Jewel Seam for unexplored parts of the Cadomin-Luscar coalfield.

#### PETROGRAPHIC CHARACTERISTICS AND TECHNOLOGICAL PROPERTIES

Technological properties of coals are to a large extent a direct function of rank and composition of the parent coal. While no special requirements exist in terms of rank and composition for thermal coals used for generation of electricity, coking coals used in carbonization to produce metallurgical coke are restricted to a fairly narrow rank range, that is from high volatile A coal ( $R_{max} > 0.95\%$ ) to the transition from medium to low volatile bituminous coals ( $R_{max} < 1.50\%$ , Jasienko (1978)). The same is also true for conversion processes such as coal liquefaction, where optimum conversion rates are obtained using coals of high volatile bituminous rank (Whitehurst et al., 1980). Within the optimum rank ranges the maceral composition determines largely the nature and quality of the final products.

Technologically, the macerals of coals are generally classified as reactive or inert. The reactive group comprises macerals of the vitrinite and liptinite groups and a portion of the maceral semifusinite

(30 - 50%). This group of macerals when exposed to processes such as carbonization will react in a sense that the organic material, depending on rank and experimental conditions used, will give rise to a variety of newly formed materials, i.e. liquids, gaseous products and residual material (cokes). The inert constituents are comprised of the macerals of the inertinite group including 50 - 75% of the maceral semifusinite. In terms of reflectance ranges, all samples from the Jewel Seam collected in the Luscar-Cadomin area fall well within the limits considered to be suitable for coke making (0.95 - 1.50% R<sub>max</sub>). Within this range the ratios of inert/reactive components will then define the quality of the final product (coke). There exists a number of empirical methods for the prediction of coke quality from petrographic analyses (Stach, 1982).

The petrographic data from 15 samples of the Jewel Seam have been plotted in a diagram modified from Pearson (1980). This diagram uses mean maximum reflectances of vitrinite and the contents in inert macerals to estimate coke stability factors. The dashed line indicates the optimum ratio of inerts to reactive macerals for producing the strongest coke for a given rank. The line of the 50 coke stability mark indicates the threshold separating strong cokes (>50) from those less acceptable (<50). From the graph shown in figure 14 it appears that the coke stability factors predicted for cokes from the Jewel Seam are close to or exceed coke stability factors of 50. This confirms the well known fact that coke produced from the Jewel Seam is of high quality.

#### MATURATION AND HYDROCARBON POTENTIAL

The burial histories and the amount and type of organic matter in the sedimentary rocks determine the formation of oil and gas. Vitrinite reflectances determined on the coals from the Luscar area indicate that the maturation levels fall within the so called "oil window" concept, in which oil generation is believed to occur in the vitrinite reflectance




Figure 14. Prediction of coke stability factors from petrographic characteristics for Jewel Seam samples of the Cadomin-Luscar area. Data do not represent current production coal.

limits of 0.5 - 1.3% (Dow, 1977). Leckie et al. (1988) showed that Lower Cretaceous strata to the north are characterized by the predominance of terrestrial organic matter, including the marine shales. This type of organic matter (kerogen III) will form gaseous products, when exposed to greater depths of burial. In other words, burial of the Lower Cretaceous strata in the Luscar area has most likely led to the formation of gas from the terrestrial matter in the shales and coal seams, while oil was not formed because of the lack of precursor materials (kerogen types I and II).



## ROADLOGS AND STOPS

Day 1: Road log between Edson and Cadomin (Highways 47 and 40).  
Kms.

- 0 Turnoff overpass from Highway 16 onto Highway 47 - Robb Road.  
View of the Front Ranges.
- 3.0 Railroad overpass.
- 8.0 Post-glacial dune field exposed along the road for the next few kilometers.
- 13.0 McLeod River bridge.
- 18.0 Entering region of glacial lacustrine silt deposits.
- 31.5 Railroad crossing.
- 35.7 Small outcrop of Cordilleran till on right.
- 36.1 View of Nikanassin Range.
- 40.0 Roadcut exposing tills, sands and volcanic ash - possibly glacially thrust bedrock.
- 40.1 Prest Creek.
- 47.2 Cordilleran till outcrop.
- 54.3 Paskapoo Formation outcrops along the highway.
- 57.3 Glacial outwash sands with fine coal debris.
- 57.4 Second turnoff to Robb townsite.
- 58.0 Northeast dipping Paskapoo Formation.
- 59.6 Northeast dipping Brazeau Formation outcrop. Palynological work by the GSC indicates an Early to Middle Maastrichtian age.
- 60.6 Axial trace of Coalspur Anticline exposed in Brazeau Formation roadcut to the right (north). Strata is roughly equivalent to the Plains Bearpaw Formation. However, the whole Brazeau Formation is continental in this area.
- 60.9 West dipping Brazeau Formation - west limb of Coalspur Anticline.
- 61.2 Embarras River - Brazeau Formation.
- 62.8 Excellent exposure of the Coalspur Formation (Scollard equivalent) and a burning coal seam (Val d'Or Seam). GSC palynological dating suggests an early Paleocene age, similar in age to the Ardley coal zone. The Cretaceous-Tertiary boundary is situated near the base of the Mynheer Seam, which is lower in the stratigraphy, but not exposed in this section.
- 63.4 Highway takes a sharp turn to the left (southeast). Good view of abandoned coal mine on hillside above old town of Coalspur.
- 64.9 Turnoff to Cadomin via Highway 40. Embarras river and outcrops of Paskapoo Formation.
- 70.5 View of the Nikanassin Range. Luscar Mountain (left), Mount Gregg (right).
- 72.5 Road to Mercoal village. Mercoal was the first opened and the last mining village to be abandoned in the Coal Branch area (1959).
- 73.1 Roadcut of east dipping Entrance Conglomerate (roughly equivalent to the Battle Formation in the Plains).
- 74.8 Railroad overpass - Steeper siding.
- 75.7 Roadcut down into the McLeod River valley exposing Brazeau Formation.

- 78.2 View of Nikanassin Range with Cadomin Mountain.
- 81.4 Highway travels along the McLeod River valley, gravel outcrops are from associated terrace deposits.
- 84.2 Nikanassin Range. Leyland Mountain on left, Luscar Mountain - center.
- 85.6 McLeod River crossing, Brazeau Formation outcrops to the right along the river.
- 87.7 Brazeau Formation on left by railroad tracks.
- 90.0 Outcrops along Watson Creek, Brazeau Formation.
- 90.4 Watson Creek campground turnoff.
- 91.4 McLeod River bridge. At Hell's Gate (1 km upstream) contact between Wapiabi and Brazeau Formations. Leyland Mountain straight ahead.
- 92.4 View of Inland Cements limestone quarry on Cadomin Mountain. Limestone is quarried from the Devonian Palliser Formation near the upper benches and is then dropped through a vertical shaft to a crusher located within the mountain. Crushed rock is then moved by belt conveyor, horizontally, out of the mountain to a rail load-out facility.
- 94.0 Cardium Formation outcrops on the right (north) side of the highway. We are now on the map of figure 15.
- 94.8 Junction with Forestry Trunk road to Cadomin. Proceed to Cadomin.
- 96.9 Blackstone Formation, with the subsurface "second white specs" horizon, exposed almost continuously to the left along the railway.
- 98.2 Blackstone Formation with subsurface "Fish Scale horizon" exposed along railway tracks to the left.
- 98.3 Greenish weathering Mountain Park Formation exposed on the right. Approximately equivalent to the Harmon or Notikewin Members of the Deep Basin area.
- 99.1 Across the river to the left is an outcrop exposure of the Lower Cretaceous (Aptian) Gladstone to Moosebar, continental to marine transition that will be examined at Stop 1. This sequence is approximately equivalent to the McMurray to Clearwater transition in northeastern Alberta and the Gething to Wilrich transition in the northern Deep Basin.
- 100.4 Approximate trace of the Nikanassin thrust, which has brought Devonian Palliser and Mississippian Rundle Group strata on top of Jurassic to Cretaceous age Fernie and Nikanassin Formations.
- 101.7 Turnoff to Inland Cement quarry and bridge over McLeod River.
- 102.0 Road runs northward along CNR tracks, exposing progressively younger strata in the southwest limb of the Cadomin Syncline (figure 15).
- 102.4 Thinly and rhythmically bedded Nikanassin Formation. In southern Alberta the correlative strata, Kootenay Group, is coal-bearing and consists of coastal plain to alluvial plain deposits.

- 102.5 Road passes over the Cadomin Formation (figure 15), which is a reliable mapping unit in this area. The Cadomin is thought to represent a series of alluvial fan to alluvial plain deposits that flowed, primarily, to the northeast from here towards the Boreal Sea. The base of the Cadomin represents a major unconformity and perhaps a major sequence boundary between the Fernie/Nikanassin and Luscar/Mannville clastic wedge sequences.
- 103.0 Road passes by folded and faulted strata of the Gladstone Formation. The section of Stop 1 was measured along these exposures.
- 103.2 Old mine entrance in Jewel Seam above Torrens sandstone.




Figure 15. Geological map of the Cadomin area, modified from MacKay (1929).

STOP 1. CONTINENTAL TO MARINE TRANSITION - GLADSTONE TO MOOSEBAR FORMATIONS.

Objective: The objective of this stop is to demonstrate the transition from continental alluvial plain strata of the Gladstone into the fully marine Moosebar Formation (figures 15 and 16).

Background: This section has been studied by several authors and microfossils have been collected from this section (McLean and Wall, 1981, Mellon, 1967). Some debate exists as to where to place the contact between the Gladstone and Moosebar Formations at this location. Examination of the section will show the difficulty, and perhaps, somewhat arbitrary nature of any pick chosen.

Things to see:

- 1). In the lower portion of the section (between 12 and 24 m) note the three lensoid shaped channel deposits. These channels tend to be small in size, are ripple laminated and contain abrupt abandonment overbank phases. Thin coals and carbonaceous mudstones are common (figure 18).
- 2). Up to the 85 m level, note the abundant indicators of continental subaerial sedimentation in this part of the succession (rootlets, logs, paleosols, plant debris, possible dinosaur tracks - figure 19). Microfossils support a fully continental interpretation for this interval.
- 3) Between 89 and 100 m note the major upward-coarsening sequence, fewer continental indicators, and sudden presence of trace fossils. Brackish conditions are inferred from microfossils collected in this interval. The upward-coarsening cycle is thought to be the Bluesky Member, which produces gas near Edson.
- 4) At 108 m note the small fault that has repeated at least one of the upward-coarsening cycles above the Bluesky (figure 20).
- 5) Above 108 m and till the end of the section, note the paucity of plant, root and other continental indicators and note the invertebrate fossil bed. Fully marine conditions have been inferred from foraminifera collected from this interval (McLean and Wall, 1981). Mellon (1967) and Mclean and Wall (1981) placed the base of the Moosebar at the invertebrate fossil bed. We would prefer to place it at the top of the Bluesky (figure 17), which is 2.5 m below the fossil bed and above the highest sandstone bed. In this way the contact will conform to a major response on geophysical logs.




Figure 16. Stratigraphic section showing the Gladstone to Moosebar Formations transition, Cadomin Railway section, Stop 1.




Figure 17. Gladstone and Moosebar formations, transition from alluvial plain to marine environments, Stop 1. Cadomin railway section.




Figure 18. Small lens shaped fluvial sandstones in the lower Gladstone Formation. Stop 1, 12 to 24m.



Figure 19. Log imprint and dinosaur cast (?) at the base of small fluvial channel, below the Bluesky member - Stop 1, 84m level.




Figure 20. Small thrust fault repeating coarsening-up unit in the lower Moosebar Formation. Stop 1, near 108m level.

## STOP 2. STRUCTURAL STYLES IN THE LOWER CRETACEOUS.

**Objective:** To demonstrate the variety of structural styles present in the Lower Cretaceous Luscar Group and their resultant effects on coal mining.

**Background:** Open to tight folds characterize the area, showing thickened coal zones in the hinges of anticlines and synclines. The open pit coal mines are generally along these thickened hinges. The present level of erosion at the Gregg River Mine results mostly in open pits along thickened syncline hinges. On the Cardinal River property open pits are located along thickened anticline and syncline hinges.

### Things to see:

- 1) On the haul road, a little past the Gregg River crossing, we pass through the mined-out core of a coal-bearing syncline (AB pit).
- 2) CD pit shows an anticline/syncline pair with a section of Moosebar to lower Grande Cache Member exposed (figure 21). Coal is thickened along the hinge, but this feature is no longer exposed.
- 3) LM pit shows a slightly recumbent syncline, with thickened coal along the hinge.
- 4) The PQ pit is situated along the thickened hinge of a syncline (figure 22). The disharmonious folding, expressed by the thickening of the Jewel Seam and tight folding of the Torrens sandstone (syncline-anticline-syncline train), results in an "H" shaped structure. The Drinnan Thrust is well exposed, with Nikanassin thrust over Moosebar, Torrens and Jewel Seam.





Figure 21. Anticline/syncline pair in CD Pit, with thickened Jewel Seam.



Figure 22. Disharmonious folding in PQ pit. The trace of the Drinnan Thrust is shown.

Day 2: Road log from Hinton to junction with Robb Highway (47).  
Kms.

- 0.0 Turnoff from Highway 16 onto Highway 40
- 1.1 Outcrop on the left (northeast) is of the Entrance Formation. An alluvial fan or braided river environment is interpreted for the Entrance. The Entrance lies at approximately the same stratigraphic position as the Battle Formation in the Plains region and is close to the Cretaceous-Tertiary boundary.
- 3.7 Cold Creek crossing.
- 7.0 View of High Divide Ridge at 10 o'clock (northeast). This ridge is an erosional remnant of mainly Paskapoo and Coalspur Formations preserved in the core of the Entrance Syncline.
- 7.7 Outcrop of coal in the Coalspur Formation near Cold Creek. The Cretaceous/Tertiary boundary is thought to lie within coals in the lower part of the Coalspur Formation.
- 9.3 Roadcut exposure on the left of the Coalspur Formation. Note the thin coal seams and volcanic ash horizon. Strata dip northeast and occur along the southwest limb of the Entrance Syncline.
- 14.6 Wigwam Creek crossing.
- 17.2 Teepee Creek crossing.
- 18.3 Upper Cretaceous Brazeau Formation exposed on left.
- 19.8 Large scale trough cross-stratification in the Brazeau Formation on the right.
- 21.2 Descending into the Gregg River valley with a good view of the Nikanassin Range. Sphinx Mountain - straight ahead composed largely of Devonian Palliser Formation (equivalent to the Wabamun platform carbonates in the subsurface).
- 23.5 Gregg River crossing.
- 24.2 Brazeau Formation outcrop on the left.
- 28.5 Platy sandstones of the Brazeau Formation on the left.
- 28.9 Small waterfall over Brazeau Formation sandstone, on the right along the Gregg River, is reported to be an excellent trout fishing spot.
- 31.4 Wapiabi Formation mudstones and shales on the left. The Wapiabi is equivalent to that portion of the Upper Colorado above the Cardium in the subsurface.
- 33.2 Cardinal River Coals 51C5 pit exposed on the northeast limb of the Luscar Anticline.
- 34.8 Crossing the approximate trace of the 10 m thick economic Jewel coal seam (Gates Formation).
- 36.4 Gregg River crossing and Jurassic-Cretaceous Nikanassin Formation exposed on left.
- 38.4 Nikanassin Formation exposed in the core of the Luscar Anticline on large outcrop to the left.
- 39.1 Gregg River Resources Limited minesite. Storage silo and load out facility visible from highway.

- 40.4 Cardinal River Coals Limited 51B2 pit (Stop 5). Mining is completed at this pit and waste rock from a developing pit to the east is now being backfilled into this pit. Pit shows the Torrens Member (sandstone) on the extreme right, overlain by the Jewel coal seam, in turn overlain by the Grande Cache Member. The thick sandstone near the middle to top is the Mountain Park Formation.
- 41.8 Bedding plane view of Torrens Member, footwall sandstone, steeply dipping to the northeast, right side of road.
- 42.4 Turnoff to Cardinal River Coals Limited mine.
- 43.9 Spoil pile being reclaimed by Cardinal River Coals.
- 44.3 From here to the Cadomin turnoff, there are several roadcut exposures of the marine Blackstone Formation (Lower Colorado/Shaftesbury equivalents).

STOP 3. THE MARINE TO NONMARINE, MOOSEBAR TO GATES TRANSITION  
(CARDINAL RIVER, 50A5 PIT).

Objective: To demonstrate the marine to nonmarine transition from the Moosebar to Gates Formation (figure 23). Show the similarities and differences with the nonmarine to marine transition seen at Stop 1. To also examine a well exposed asymmetrical anticline.

Background: This section shows a fully marine base, and passes upward into a series of upward-coarsening cycles - often capped by carbonaceous mudstones or coals. The sandstones at the top of these cycles are approximate stratigraphic equivalents of sandstones within the Wilrich A and B and the Falher G/H cycles of the northern Deep Basin area. These sandstones are gas-bearing in the northern Deep Basin and are stacked vertically. The economic Jewel Seam caps the marine sequence and is itself locally overlain by brackish water sediments. Wholly continental alluvial plain strata characterize the section up to the R Seam and likely even higher up to the top of the Mountain Park Formation.

No gas production has come from these upward-coarsening sequences in this area as yet. Why is this? The sandstones are generally very thick, well sorted and contain at least thin mudstone seals. Gas is present in this area with production coming from the Turner Valley (Mississippian), Cadomin and Bluesky. To their detriment as a potential reservoir, the sands are fine to very fine-grained and have clay or quartz overgrowth cements. Coal seams in the Elsworth Deep Basin have been examined as an important source rock for gas generation and as a potential reservoir in themselves.

Things to see:

- 1) The lowermost cycle shows bioturbation, wavy-bedding, a small fold, upright tree stump and is capped by carbonaceous shale (figure 24). At 27 m lies an upright tree stump.
- 2) A glauconitic bed lies at 33 m and may represent, in part, a transgressive condensed reworked shelf deposit. Look for bioturbation, graded-bedding, glauconite, and a pebble bed.
- 3) The second marine cycle (figure 25) is characterized by offshore mudstones at the base and passing upwards into HCS storm units and finally into parallel laminated foreshore deposits. The top of this cycle is at 75 m, and is marked by the Torrens A coal seam.
- 4) The third cycle, Torrens Member proper, consists of massive to parallel laminated, bioturbated sandstone. Look for internal scour surfaces, trough cross-stratification, HCS and roots below the Jewel. Note the very tight reservoir potential.

- 5) Jewel coal seam showing approximate original stratigraphic thickness (figure 26). Look for partings in coal, intensity of shearing, and scour at the top of the seam.
- 6) Strata above the Jewel consist of small fluvial channel and crevasse splay deposits - sometimes capped with coal and paleosols (figure 26).
- 7) The large asymmetrical anticline is well exposed in the center of this pit and consists of the Torrens Member (footwall) sandstone. The northwest plunge of the structure is clearly displayed (figure 27)






Figure 23. Stratigraphic section showing the Moosebar to Gates Formation transition, Cadomin area, Cardinal River Coals Ltd. 50A5 pit, Stop 3.




**Figure 24.** Lowermost coarsening-up cycle with overlying carbonaceous mudstones. Stratigraphic top is to the right. 50A5 pit. Stop 3.



**Figure 25.** Second coarsening-up cycle, with glauconite beds near the base. Stratigraphic top is to the right. 50A5 pit. Stop 3.



**Figure 26.** Jewel seam and overlying strata. Beds overturned with stratigraphic top to the right. 50A5 pit. Stop 3.



**Figure 27.** Well exposed anticline in the Torrens member, Stop 3, 50A5 pit. The plunge is indicated. Trucks at lower right for scale.

STOP 4. STRUCTURAL/STRATIGRAPHIC OVERVIEW AND DETAILED EXAMINATION OF THE TORRENS MEMBER (CARDINAL RIVER COAL 51B2 and 51B3 PITS).

Objective: To provide an overview of the structural and stratigraphic framework of the Grande Cache Member and to examine a problematic section of the Torrens Member.

Background: The Cardinal River Coals 51B2 and 51B3 pits have collectively exposed almost 2 km. of Grande Cache Member strata, perpendicular to the regional fold axis. The disharmonious folding style is plainly visible from the ridge as is the relationship between thickened coal and structure.

Things to see:

- 1) Thickened Jewel Seam above Torrens Member (figure 28).
- 2) Cylindrical folding of Torrens sandstone (figure 28).
- 3) Disharmonious folding as expressed by open folding of Torrens sandstone and tighter folding of strata above the Jewel Seam (figure 28 and 29).
- 4) Hinge collapse and resulting faulting (figure 28).
- 5) Roll-over folding in Torrens sandstone (figure 29).
- 6) Box fold in Rider Seam (figure 29).
- 7) Step faults in Torrens sandstone (figure 28).
- 8) Thrust fault bringing up the Jewel Seam (figure 29).
- 9) Several other faults, such as upthrusts, back thrusts and normal faults, which are probably related to the general folding (figure 29).
- 10) Combination of channel scour and thrusting (figure 29).




Figure 28. Anticline of 51B3 Pit, with disharmonious folding of Jewel Seam and nearby strata.






Figure 29. Vertical down-plunge cross section, perpendicular to the fold axis trend of N125E, through the end wall of the 51B3 Pit. For methodology of down-plunge cross section construction see Langenberg et al. (1987). TT=Top Torrens; J=Jewel Seam; GC=Grande Cache Member; R=R Seam. Sense of fault movement is indicated with arrows.

## STOP 4. continued.

Background: The second portion of the Stop will examine a problematic section of the Torrens Member that shows good evidence of some kind of channel deposition as opposed to the near-shore marine type setting usually envisaged for the Torrens (figure 30). This section is believed to represent deposition in a tidal or distributary channel setting. The reservoir potential in the subsurface of this facies of the Torrens is believed to be considerably better than the more typical sections seen, due mainly to the larger grain size and probably greater permeability.

## Things to see:

- 1) Trough cross-stratification and coarser grain size than was seen at the Stop 3 section of Torrens (from 0 to 5 m).
- 2) Breccia clasts and erosive scour surface at 6 m (figure 31).
- 3) Pebble beds, intraclasts and bedding plane view at 8 m.
- 4) Bedding plane view of large mega-ripples or dunes with overlying pebble beds at 12 m (figure 32).
- 5) Low-angle cross-stratification in very fine sandstones, more typical of the Torrens Member.






Figure 30. Stratigraphic section of a problematic Torrens Member section. Cadomin area, Cardinal River Coals Ltd., 51B2 pit. Stop 4.



**Figure 31.** Channel margin (?) and breccia beds in an atypical Torrens section. Stop 4, 51B2 pit vicinity.



**Figure 32.** Mega-ripples exposed on bedding plane surface, near the middle of an atypical Torrens section. Stop 4, 51B2 pit vicinity.

STOP 5. COASTAL PLAIN TO ALLUVIAL PLAIN ENVIRONMENTS OF THE GRANDE CACHE MEMBER (CARDINAL RIVER, 51B2 PIT).

Objective: To examine the coastal plain to alluvial plain environments in the Grande Cache Member, above the Jewel coal seam (figure 33). Compare the reservoir potential of these sandstones to those in the cyclic marine upward-coarsening sandstones below the Jewel Seam.

Background: This portion of the section will examine the coastal to alluvial plain transition above the Jewel Seam at the Cardinal River Coals 51B2 pit. Figure 9 shows that this transition can be broken up into 5 distinct lithofacies units:


- 1) coastal mire (GC1),
- 2) Tidal or brackish (GC2),
- 3) amalgamated channels (GC3),
- 4) small fluvial channels (GC4) and
- 5) floodplain (GC5).


Things to see:

- 1) In the first 8.5 m above the Jewel Seam (GC2 facies), look for indications of brackish water deposition (tidal bedding). Forams have been recovered from this interval.
- 2) The next 25 m (up to the 38 m level) shows amalgamated channels (GC3 facies) with internal scour surfaces, log impressions, trough cross-stratification, lateral accretion surfaces and a slump block. Medium grain-size and thickness may make this an excellent subsurface reservoir (figure 34).
- 3) At 36 m is the base of the GC4 lithofacies unit. This unit has several stacked, small-scale fluvial channels, most of which fine upwards. Look for ripple lamination, fining-up cycles and thin carbonaceous mudstones (up to the 50 m level).
- 4) The floodplain facies (GC5) in this area is anomalous in that thin fluvial channel deposits predominate over mudstones. Look for fining-up cycles, predominance of low energy current indicators, roots below coals and rhythmic bedding near the R coal seam. Note the multiple partings and poor quality of the R Seam (at the 80 m level).
- 5) An upright tree stump is preserved about 8 m above the R Seam (figure 35).
- 6) Thrust visible from the road, which places Moosebar on top of Torrens (figure 36)


Figure 33. Stratigraphic section of the coastal plain to alluvial plain transition, Jewel coal to middle Grande Cache member, 51B2 pit, Stop 5.

Section 4






**Figure 34.** Amalgamated channel facies (GC3) showing lateral accretion surfaces. Photo rotated to correct for dip. Stop 5, 51B2 pit. Sandstone about 10m thick.



**Figure 35.** Upright tree stump, 8m above R seam. Stop 5, 51B2 pit.



**Figure 36.** Thrust emplacing Moosebar Formation on top of the Torrens Member. Stop 5, 51B2 pit.

STOP 6. GATES FORMATION (MOUNTAIN PARK MEMBER) TO BLACKSTONE FORMATION TRANSITION AND ABSENCE OF FOLDING MOUNTAIN FAULT.

Objective: The objective of this stop will be to observe the fluvial/alluvial plain Mountain Park Member to fully marine Blackstone Formation transition.

Background: The Blackstone transgression in this area appears to be very abrupt. A thin transgressive pebble lag deposit is the only evidence remaining of the transgression itself. This does not necessarily imply that the transgression was in itself a rapid event - it may simply be that most of the transgressive associated sediments have been reworked and simply not been preserved in this area. This contact is probably a major sequence boundary. Contrast this alluvial to marine transition with the Gladstone to Moosebar transition seen at Stop 1.

According to mapping by Mountjoy (1959) and MacKay (1929) a major thrust fault (the Folding Mountain Thrust) is thrusting the Mountain Park Member on top of the Blackstone Formation. However, no evidence of faulting is found in this section. Faulting may be present in the Blackstone Formation (Blackstone Zone of Detachment), but the orientation of this zone is different from that of the postulated Folding Mountain Thrust.

Things to see:

- 1) Greenish colored fluvial sandstones of the Mountain Park Formation.
- 2) Thin transgressive chert pebble bed at the Mountain Park/Blackstone contact.
- 3) Fully marine dark grey mudstones with thin sandstones in the Blackstone Formation.
- 4) The absence of a major thrust fault, indicated by the normal stratigraphic contact of the Mountain Park Member and the Blackstone Formation.



## REFERENCES

- Bustin, R.M., Cameron, A.R., Grieve, D.A. and Kalkreuth, W.D. (1983):  
Coal petrology, its principles, methods and applications.  
Geological Association of Canada, Short Course Notes, v. 3,  
272 pages.
- Cant, D.J. (1983): Spirit River Formation—A stratigraphic diagenetic gas  
trap in the deep basin of Alberta; Bulletin American Association of  
Petroleum Geologists, v. 67, pp. 577-587.
- Cecil, C.B., Stanton, R.W., Dulong, F.T. and Renton, J.J. (1980):  
Geologic factors that control mineral matter in coal.  
*In* Donaldson, A.C., Presley, M.W. and Renton, J.J. (editors),  
Carboniferous coal short course and guidebook, v. 3, American  
Association of Petroleum Geologists field seminar, 1980. pp. 43-56.
- Diessel, C. (1982): An appraisal of coal facies based on maceral  
characteristics. Australian Coal Geology, v. 4, Part 2, pp.  
131-146.
- Diessel, C. (1986): The correlation between coal facies and depositional  
environments. Advances in the study of the Sydney Basin, Proc. of  
the 20th Symposium, The University of Newcastle, pp. 19-22.
- Dow, W. (1977): Kerogen studies and geological interpretations. Journal  
of Geochemical Exploration, v. 7, pp.79-99.
- ERCB, (1987): Reserves of coal, Province of Alberta, ERCB Report  
ST88-31.
- Englund, K.J., Windolph, J.F. and Thomas, R.E. (1986): Origin of thick,  
low sulfur coal in the Lower Pennsylvanian Pocahontas Formation,  
Virginia and West Virginia. Geological Society of America Special  
Paper 210, pp. 49-61.

- Ekdale, A.A., Bromley, R.G. and Pemberton, S.G. (1984): Ichnology - The use of trace fossils in sedimentology and stratigraphy, Society of Economic Paleontologist and Mineralogists, Short course notes no. 5, 317p.
- Gies, R.M., (1984): Case history for a major Alberta Deep Basin gas trap: the Cadomin Formation. Edited by Masters, J. American Association of Petroleum Geologists, Memoir 38, pp. 115-140.
- Haq, B.U., Hardenbol, J. and Vail, P.R. (1987): Chronology of fluctuating sea levels since the Triassic. *Science*, v. 235, no. 6, pp. 1156 - 1167.
- Hill, K.C. (1980): Structure and stratigraphy of coal-bearing strata near Cadomin, Alberta. Unpublished M.Sc. thesis, University of Alberta, 191 p.
- Jackson, P.C. (1984). Paleogeography of the Lower Cretaceous Mannville Group of Western Canada. *In* Elmworth - Case study of a deep basin gas field. Edited by Masters, J. American Association of Petroleum Geologists, Memoir 38, pp.79-114.
- Jasienko, S. (1978): The nature of coking coals. *Fuel*, v. 57, pp. 131-146.
- Kalkreuth, W. (1988): Kanadische Kreidekohlen- Inkohlungsgrad und petrographische Zusammensetzung Kanadischer Kreidekohlen in den Rocky Mountain Foothills von NE-British Columbia und Westzentral Alberta. *Erdol und Kohle*, v. 41, p. 2, pp. 61-70.
- Kalkreuth, W. and Leckie, D.A. (in press): Sedimentological and petrographical characteristics of Cretaceous strandplain coals: A model for the accumulation from the North American Western Interior seaway. *Journal of Coal Geology*.
- Langenberg, C.W. and McMechan, M.E. (1985): Lower Cretaceous Luscar

- Group (revised) of the northern and north central Foothills of Alberta; Bulletin of Canadian Petroleum Geology, v. 33, pp. 1-11.
- Langenberg, C.W., Kalkreuth, W. and Wrightson, C.B. (1987): Deformed Lower Cretaceous coal-bearing strata of the Grande Cache area, Alberta. Alberta Research Council, Bulletin 56, 54p.
- Langenberg, C.W., Macdonald, D.E., Kalkreuth, W., Strobl, R.S. and Bahnsen, B. (1988): An initial assesment of controls on coal quality variation in the Cadomin-Luscar coalfield. Alberta Research Council, Open File Report 1988-06.
- Leckie, D.A. and Walker, R.G. (1982): Storm- and tide- dominated shorelines in the Cretaceous Moosebar - Lower Gates interval - outcrop equivalents of Deep Basin gas trap in Western Canada. American Association of Petroleum Geologist Bulletin, v. 66, pp. 138-157.
- Leckie, D.A., Kalkreuth, W.D., and Snowden, L.R. (1988): Source rock potential and thermal maturity of Lower Cretaceous strata: Monkman Pass area, British Columbia. American Association of Petroleum Geologist Bulletin, v. 72, pp. 1820-838.
- Macdonald, D.E., Langenberg, C.W. and Strobl, R.S. (1988): Cyclic marine sedimentation in the Lower Cretaceous Luscar Group and Spirit River Formation of the Alberta Foothills and Deep Basin. *In* Sequences, Stratigraphy, Sedimentology: Surface and Subsurface, eds. D. P. James and D. A. Leckie, Canadian Society of Petroleum Geologists, Memoir 15, pp. 143-154.
- MacKay, B.R. (1929): Cadomin, Alberta. Geological Survey of Canada, Map 209A.
- MacKay, B.R. (1930): Stratigraphy and structure of the bituminous coalfields in the vicinity of Jasper Park, Alberta. CIMM, Bulletin No. 222, Transactions Section, pp. 1306-1342.



















































- McCabe, P.J. (1984): Depositional environments of coal and coal-bearing strata. Special Publication of the International Association of Sedimentologist, no. 7, pp. 13-42.
- McCabe, P.J. (1987): Facies studies of coal-bearing strata, from Scott, A.C. (eds), Coal and coal-bearing strata: recent advances, Geological Society of Special Publication no. 32, pp. 51-66.
- McLean, J.R. (1977): The Cadomin Formation: stratigraphy, sedimentology and tectonic implications; Bulletin of Canadian Petroleum Geology, v. 25, pp. 792-827.
- McLean, J.R. (1982): Lithostratigraphy of the Lower Cretaceous coal-bearing sequence, Foothills of Alberta; Geological Survey of Canada, Paper 80-29, 46p.
- \_\_\_\_\_ and Wall, J.H. (1981): The early Cretaceous Moosebar Sea in Alberta; Bulletin of Canadian Petroleum Geology, v. 29, pp. 334-377.
- Mellon, G.B. (1966): Lower Cretaceous section, Cadomin area, Alberta. *In* Williams, G.D. (editor), Edmonton Geological Society Eighth Annual Field Trip Guidebook to Cadomin, August, 1966.
- Mellon, G.B. (1967): Stratigraphy and petrology of the Lower Cretaceous Blairmore and Mannville groups, Alberta Foothills and Plains. Alberta Research Council, Bulletin 21, 270 pages.
- Monger, J.W.H. (1984): Cordilleran tectonics: Canadian perspective. Bulletin de la Societe Geologique de France, v. 26, pp. 255-278.
- Mountjoy, E.W. (1959): Miette, Alberta. Geological Survey of Canada, Map 40-1959.
- Pearson, D. (1980): The quality of Canadian coking coal. CIMM Bulletin, v. 73, no. 813, pp. 70-84.

- Smith, D.G., Zorn, C.E. and Sneider, R.M. (1984): The paleogeography of the Lower Cretaceous of western Alberta and northeastern British Columbia in and adjacent to the Deep Basin of the Elmore area; *in* Elmore - Case study of a deep basin gas field. Edited by Masters, J.A., Journal American Association of Petroleum Geologists, Memoir 38, pp. 79-114.
- Stach, E., Mackowsky, M.-Th., Teichmüller, M., Taylor, G., Chandra, D., Teichmüller, R. (1982): Stach's Textbook of Coal Petrology, 2nd. edition, Gebrüder Borntraeger, Stuttgart, 428p.
- Stott, D.F. (1984). Cretaceous sequences of the Foothills of the Canadian Rocky Mountains, Alberta; *in* The Mesozoic of Middle North America, edited by D.F. Stott and D.J. Glass: Canadian Society of Petroleum Geologists, Memoir 9, pp. 85-107.
- Strobl, R.S. (1988): The effects of sea-level fluctuations on prograding shorelines and estuarine valley-fill sequences in the Glauconitic Member, Medicine River Field and adjacent areas. *In*: Sequences, Stratigraphy, Sedimentology: Surface and Subsurface, eds. D. P. James and D. A. Leckie, Canadian Society of Petroleum Geologists, Memoir 15, pp. 221-236.
- Taylor, D.R. and Walker, R.G. (1981): Depositional environments and paleogeography in the Albian Moosebar Formation and adjacent fluvial Gladstone and Beaver Mines Formations, Alberta; Canadian Journal of Earth Sciences, v. 21, p. 698-714.
- Walker, R.D. (1984): Shelf and shallow marine sands. *In* Facies Models. Edited by Walker, R.D., Geological Association of Canada, Reprint Series 1, pp. 141-170.

- Weimer, R.J. (1983): Relation of unconformities, tectonics, and sea level changes, Cretaceous of the Denver Basin and adjacent areas. *In* Mesozoic paleogeography of west - central United States, eds. Reynolds, M. W. and Dolly, E.D., Society of Economic Paleontologist and Mineralogists, Rocky Mountain Section, pp. 359-376.
- Whitehurst, D., Mitchell, T. and Farcasiu, M. (1980): Coal liquefaction. Academic Press, New York, 378p.

APPENDIX

Legend:

	Sandstone		Trough x-strata		Heavily rooted
	Conglomerate		Ripples		Soil mottled
	Interbedded mudstone/coal (<25% coal)		Soft sediment deformation		Symmetrical ripples
	Clay-shale Claystone		Roots		Wavy bedding
	Siltstone Silshale		Leaf imprints		
	Interbedded claystone Mudstone (<25% claystone)		Log or stems		
	Mudstone Mudshale		Basal lag		
	Interbedded sandstone Claystone (>50% sandstone)		Parallel stratification		
	Interbedded sandstone Claystone (>75% sandstone)		Intraclasts		
	Paleosol		Graded bedding		
	Coal		Slickensides		
	Low-angle x-strata		Folding		
	Hummocky cross stratification		Fault		
	Very fn interlaminated		Glaucconitic		
	Carbonaceous matter-particles		Pyrite		
	Shells (marine)		Bioturbated		
	Fining-upward cycle		Forams		
	Coarsening-upward cycle		Gradational contact		
	Carbonaceous matter-finely disseminated		Sharp contact		
	Carbonaceous matter in thin laminae		Burrows (vertical horizonta, with spriten, branching)		
	Scour and fill		Ironstone band		
	Sheared coal		Erosional contact		
	Large scale x-bedding		Dinosaur track		